



Education, conservation, and research from the world's first cetacean protection organization.

 Search

Gray Whale

Eschrichtius robustus

CLASS: Mammalia

ORDER: Cetacea

SUBORDER: Mysticeti

FAMILY: Eschrichtiidae

GENUS: Eschrichtius

SPECIES: robustus




The only member of the family Eschrichtiidae, the gray whale is a mysticete, or baleen whale. It is a "coastal" whale that migrates along the North American Pacific Coast between arctic seas and the lagoons of Baja California, Mexico. Frequently visible from shore, gray whales provide a unique opportunity for land and boat observation, and commercial whale watching has become a major industry along its migration route. Visitors to the calving and breeding lagoons sometimes encounter the phenomenon of the "friendlies"; gray whales that closely approach small boats and allow themselves to be touched by humans.

PHYSICAL DESCRIPTION

Gray whales have a streamlined body, with a narrow, tapered head. The upper jaw is arched in profile, and slightly overlaps the lower jaw. The rostrum (upper jaw) is dimpled and each of the little depressions contains one stiff hair. There are 2-5 grooves on the ventral throat 5 feet (1.5 m) in length.

COLOR

The gray whale received its name from the gray patches and white mottling on its dark skin. On the skin are many scratches, scattered patches of white barnacles, and orange whale lice. Newborn calves are dark gray to black, although some may have distinctive white markings.



Surface Characteristics

FINS AND FLUKE

The gray whale has no dorsal (top) fin. About 2/3 of the way back on its body is a prominent dorsal hump followed by a series of 6-12 knuckles along the dorsal ridge that extend to the flukes (tail lobes). Its flippers are paddle shaped and pointed at the tips. Its fluke is about 10-12 feet (3.7 m) across, pointed at the tips, and deeply notched in the center.

LENGTH AND WEIGHT

Adult males measure 45-46 feet (13.7-14 m) and adult females measure slightly more. Both sexes weigh 30-40 tons (27,200-36,300 kg).

FEEDING

Gray whales feed on small crustaceans such as amphipods, and tube worms found in bottom sediments. They feed primarily during the summer months of long daylight hours in the cold Arctic waters of the Bering and Chukchi seas. As a baleen whale, it has a series of 130-180 fringed overlapping plates hanging from each side of the upper jaw, where teeth might otherwise be located. These plates consist of a fingernail-like material called keratin that frays out into fine hairs on the ends inside the mouth next to the tongue. The plates are off-white and about 2-10 inches (5-25 cm) in length. To feed, a whale dives to the bottom, rolls on its side and draws bottom sediments and water into its mouth. As it closes its mouth, water and sediments are expelled


through the baleen plates, which trap the food on the inside near the tongue to be swallowed.

MATING AND BREEDING

Gray whales reach sexual maturity at 5-11 years of age, or when they reach 36-39 feet (11-12 m) in length. Gestation is 12-13 months. The calf weighs 1,100-1,500 pounds (500-680 kg) and is about 15 feet (4.5 m) at birth. Calves nurse 7-8 months on milk that is 53% fat (human milk is 2% fat). Females bear a single calf, at intervals of 2 or more years. Courtship and mating behavior are complex, and frequently involve 3 or more whales of mixed sexes. Mating and calving both occur primarily in the lagoons of Baja California, Mexico, although both have been observed during the migration.

DISTRIBUTION AND MIGRATION

Gray whales inhabit shallow coastal waters of the eastern North Pacific. The gray whale makes one of the longest of all mammalian migrations, averaging 10,000-14,000 miles (16,000-22,530 km) round trip. In October, the whales begin to leave their feeding grounds in the Bering and Chukchi Seas and head south for their mating and calving lagoons in Baja California, Mexico. The southward journey takes 2-3 months. The whales remain in the lagoons for 2-3 months, allowing the calves to build up a thick layer of blubber to sustain them during the northward migration and keep them warm in the colder waters. The return trip north takes another 2-3 months. Mothers and calves travel very near shore on the northbound migration. There are some individual gray whales that are found year round in the Straits of Juan de Fuca between the State of Washington and Vancouver Island, Canada, and some that are seen during the summer months off the northern California coast.



Range Map

NATURAL HISTORY

A migrating gray whale has a predictable breathing pattern, generally blowing 3-5 times in 15-30 second intervals before raising its fluke and submerging for 3-5 minutes. A gray whale can stay submerged up to 15 minutes, and travel at 3-6 miles per hour (4.8-9.6 km/hr). Mothers are very protective of their calves, and earned the name "Devilfish" from early whalers in the lagoons because of their violent defensive behaviors. Orcas (killer whales) are a cause of gray whale deaths, and many gray whales have orca teeth scars on their flukes.

STATUS

At one time there were three gray whale populations: a north Atlantic population, now extinct, possibly the victims of over-hunting; a Korean or western north Pacific stock now very depleted, also possibly from over-hunting; and the eastern north Pacific population, the largest surviving population. Hunted to the edge of extinction in the 1850's after the discovery of the calving lagoons, and again in the early 1900's with the introduction of floating factories, the gray whale was given partial protection in 1937 and full protection in 1947 by the International Whaling Commission (IWC). Since that time the eastern north Pacific gray whale population has made a remarkable recovery and now numbers between 19,000 and 23,000, probably close to their original population size.

BIBLIOGRAPHY

Gohier, F. 1988. A Pod of Gray Whales. Blake Publishing, San Luis Obispo, California. Jones, M.L. and S.L. Swartz. 2002. Gray Whale, *Eschrichtius robustus*. In: Encyclopedia of Marine Mammals. Perrin, W.F., B. Wursig, and J.G.M. Thewissen (eds). Academic Press, San Diego. Pp. 524-536. Jones, M. L., Swartz, S. L., and Leatherwood, S., eds. 1984. The Gray Whale. Academic Press, New York, NY. Reeves, R.R., Stewart, B.S., Clapham, P.J., and Powell, J.A. 2002. Guide to Marine Mammals of the World. Alfred A. Knopf, New York.

ACKNOWLEDGEMENTS:

We greatly appreciate the knowledge and assistance of Steven Swartz of the United States National Marine Fisheries Service, who contributed to the revision of this fact sheet. Illustrations courtesy Uko Gorter, copyright© 2003, 2006 all rights reserved.

FACT SHEETS MAY BE REPRINTED FOR EDUCATIONAL OR SCIENTIFIC PURPOSES

American Cetacean Society

P.O. Box 1391, San Pedro, CA 90733-1391
310-548-6279 (Office) • 310-548-6950 (fax) • acsoffice@acsonline.org